UAP

Partition

DEPOSITION AID AND DRIFT RETARDANT

PRINCIPLE FUNCTIONING AGENTS:

~ ~	0.7
30	%
70	%
100	0/-
100	/0
	30 70 100

All ingredients are accepted for use under 40 CFR 180.1001(d).

CAUTION

KEEP OUT OF REACH OF CHILDREN

May cause irritation to skin and eyes. If in contact flush with plenty of water, see a doctor if irritation persists.

NOTE: Read and follow precautions on label of pesticide used.

STATEMENT OF PRACTICAL TREATMENT

Eye Contact: Immediately flush with water for 15 to 20 minutes.

Seek medical attention if irritation persists.

Skin Contact: Wash with soap and water. Seek medical

attention if irritation persists.

Inhalation: Remove to fresh air, treat symptomatically.

Seek medical attention if irritation persists.

If Ingested: Rinse out mouth and drink 2 glasses of water.

DO NOT INDUCE VOMITING. Seek immediate

medical attention.

GENERAL INFORMATION

Partition is an effective, easy-to-use adjuvant, for deposition improvement and drift retardation in spraying operations. **Partition** is compatible with most water-soluble and wettable powder pesticides, desiccants and cotton defoliants when applied by aerial application or standard ground equipment.

DIRECTIONS FOR USE

IMPORTANT: Keep container closed in storage and do not allow water to come in contact with contents until added to the spray solution. In case of contact with skin or eyes, flush immediately with plenty of water.

- Step 1. Select correct dosage from chart below.
- **Step 2.** Fill mix tank with water, pesticide and other additives and begin to agitate.
- Step 3. SHAKE BOTTLE OF PARTITION WELL BEFORE USING.
- **Step 4.** For best mixing results, either (1) inject **Partition** liquid into the suction side of the feeder, or recirculating pump to obtain adequate dispersion of the polymer throughout the pesticide solution, or (2) slowly add **Partition** liquid to the rapidly agitating tank mix in the area of highest turbulence by pouring a very thin stream
- **Step 5.** Continue to agitate mix tank for at least 2 minutes before spraying.
- **Step 6**. Follow all cleanup precautions on label of pesticide used. Follow governmental procedures for disposal of pesticide spray solution.

Issued: 03/04

DOSAGE CHART

SPRAY	NOZZLE	PARTITION
PRESSURE	ORIENTATION	*DOSAGE
Aerial:	Straight Back	6 - 12 oz.
below (45 psi)	45° Angle Back	8 - 16 oz.
Ground:	Flat Fan, Flooding	4 - 8 oz.
low (below 30 psi)	Off-Center	6 - 10 oz.
medium (30-50 psi)	Flat Fan, Flooding Off-Center Spray Guns	6 - 8 oz. 8 - 12 oz. 10 - 16 oz.
high (over 50 psi)	Spray Guns	16 - 64 oz.

*per 100 gallons of spray solution.

For home and garden sprayers, use approximately 1 Tablespoon of **Partition** per 5 gallons of spray solution.

STORAGE AND DISPOSAL

PROTECT FROM FREEZING. Store above 32°F. Partition's shelf life is guaranteed for six months from the ship date. Do not reuse empty container. To avoid product degradation and equipment corrosion, do not use iron, copper or aluminum containers or equipment. Dispose of container in a safe manner in accordance with federal, state and local regulations.

USE PRECAUTIONS

The degree of drift hazard varies with the type of pesticide, application conditions, and vegetation near the sprayed area. Consult your local agricultural advisor. Remember, pesticide drift is no accident. Common sense and sound application technology must be followed when spraying pesticides. **Partition** will retard, but not totally eliminate drift.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

